

Referenze

Di seguito una selezione della letteratura che testimonia l'importante contributo della Terapia Occupazionale negli ambiti di interesse del Piano Nazionale delle Cronicità.

- Adams, J. (1996). Splinting the rheumatoid wrist and hand: Evidence for its effectiveness. *British Journal of Therapy & Rehabilitation*, 3(11), 621-624.
- Adams, J., Burridge, J., Mullee, M., Hammond, A., & Cooper, C. (2008). The clinical effectiveness of static resting splints in early rheumatoid arthritis: A randomized controlled trial. *Rheumatology*, 47(10), 1548-1553.
- Adams, J., Burridge, J., Mullee, M., Hammond, A., & Cooper, C. (2004). Correlation between upper limb functional ability and structural hand impairment in an early rheumatoid population. *Clinical Rehabilitation*, 18(4), 405-413. (Level III)
- Adams, J., Burridge, J., Mullee, M., Hammond, A., & Cooper, C. (2005). The effects of early rheumatoid arthritis on dominant and non-dominant hand impairment and function. *British Journal of Hand Therapy*, 10(3-4), 93-97. (Level III)
- Ahlström, S., & Bernspång, B. (2003). Occupational performance of persons who have suffered a stroke: A followup study. *Scandinavian Journal of Occupational Therapy*, 10, 88-94. (Level III)
- Ahuya, S. S., Rege, P. V., Rege, S., & Chorghade, L. U. (2006). Study of depth perception in hemiplegics. *Indian Journal of Occupational Therapy*, 27(2), 31-36. (Level II)
- Alsaker, S., & Josephsson, S. (2003). Negotiating occupational identities while living with chronic rheumatic disease. *Scandinavian Journal of Occupational Therapy*, 10(4), 167-176.
- Arbesman, M., & Pillerito, J. M. (2008). Evidence-based perspective on the effect of automobility-related modifications on the driving ability, performance, and safety of older adults. *American Journal of Occupational Therapy*, 62, 173-186. (Systematic review)
- Arokoski, J. P. (2005). Physical therapy and rehabilitation programs in the management of hip osteoarthritis. *Europa Medicophysica*, 41(2), 155-161.
- Aronsson, B., Perk, J., Norlén, A. S., & Hedbäck, B. (2000). Resuming domestic activities after myocardial infarction: A study in female patients. *Scandinavian Journal of Occupational Therapy*, 7, 39-44. (Level III)
- Austin, J., Williams, R., Ross, L., Moseley, L., & Hutchison, S. (2005). Randomised controlled trial of cardiac rehabilitation in elderly patients with heart failure. *European Journal of Heart Failure*, 7(3 SPEC. ISS.), 411-417.
- Azad, N., Molnar, F., & Byszewski, A. (2008). Lessons learned from a multidisciplinary heart failure clinic for older women: A randomized controlled trial. *Age and Ageing*, 37, 282-287. (Level I)
- Baldelli, M. V., Pradelli, J. M., Zucchi, P., Martini, B., Orsi, F., & Fabbo, A. (2007). Occupational therapy and dementia: The experience of an Alzheimer Special Care Unit. *Archives of Gerontology and Geriatrics*, 44(Suppl. 1), 49-54. (Level III)
- Barry, M. A., Purser, J., Hazleman, R., McLean, A., & Hazleman, B. L. (1994). Effect of energy conservation and joint protection education in rheumatoid arthritis. *British Journal of Rheumatology*, 33(12), 1171-1174.
- Bendstrup, K. E., Ingemann-Jensen, J., Holm, S., & Bengtsson, B. (1997). Out-patient rehabilitation improves activities of daily living, quality of life and exercise tolerance in chronic obstructive pulmonary disease. *The European Respiratory Journal: Official Journal of the European Society for Clinical Respiratory Physiology*, 10(12), 2801-2806.

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale:

Via Angelo Emo 144, 00136 Roma

Email / sito web:

info@aito.it / www.aito.it

- Berggren, M., JoostDavidsson, A., Lindstrand, J., Nylander, G., & Povlsen, B. (2001). Reduction in the need for operation after conservative treatment of osteoarthritis of the first carpometacarpal joint: A seven year prospective study. *Scandinavian Journal of Plastic and Reconstructive Surgery and Hand Surgery / Nordisk PlastikkirurgiskForening [and] Nordisk Klubb for Handkirurgi*, 35(4), 415-417.
- Bergmann, M., Joost-Davidson, A., Lindstrand, J., Nylander, G., & Povlsen, B. (2001). Reduction in the need for operation after conservative treatment of osteoarthritis of the first carpometacarpal joint: A seven year prospective study. *Scandinavian Journal of Plastic & Reconstructive Surgery & Hand Surgery*, 35, 415-417. (Level I)
- Bilney, B., Morris, M., & Perry, A. (2003). Effectiveness of physiotherapy, occupational therapy for people with Huntington's disease: A systematic review. *American Society of Neurorehabilitation*, 17, 12-24. (Systematic review)
- Biundo, J., Jr., & Hughes, G. M. (1997). Rheumatoid arthritis rehabilitation: Part 2: Thermal modalities, occupational therapy, and psychological help. *Consultant (00107069)*, 37(12), 3123.
- Björkdahl, A., Nilsson, Å., Grimby, G., & Sunnerhagen, K. S. (2006). Does a short period of rehabilitation in the home setting facilitate functioning after stroke? A randomized controlled trial. *Clinical Rehabilitation*, 20, 1038-1049. (Level I)
- Bohr, P. C. (2000). Efficacy of office ergonomics education. *Journal of Occupational Rehabilitation*, 10, 243-255. (Level I)
- Boman, I.-L., Lindstedt, M., Hemmingsson, H., & Bartfai, A. (2004). Cognitive training in home environment. *Brain Injury*, 18, 985-995. (Level III)
- Bonnier, B., Eliasson, A.-C., & Krumlinde-Sundholm, L. (2006). Effects of constraint-induced movement therapy in adolescents with hemiplegic cerebral palsy: A day camp model. *Scandinavian Journal of Occupational Therapy*, 13(1), 13-22. (Level III)
- Borenstein, D. G., Silver, G., & Jenkins, E. (1993). Approach to initial medical treatment of rheumatoid arthritis. *Archives of Family Medicine*, 2(5), 545-551.
- Boswell, S. (1989). A social support group for depressed people. *Australian Occupational Therapy Journal*, 36(1), 34-41.
- Boter, H. (2004). Multicenter randomized controlled trial of an outreach nursing support program for recently discharged stroke patients. *Stroke*, 35, 2867-2872. (Level I)
- Bourstedt, C., Nordenskiöld, U., & Nilsson, A. L. (2009). Effects of a hand-joint protection programme with an addition of splinting and exercise. *Clinical Rheumatology*, 28, 793-799. (Level II)
- Brandis, S. J., & Tuite, A. T. (2001). Falls prevention: Partnering occupational therapy and general practitioners. *Australian Health Review*, 24(1), 37-42. (Level III)
- Branick, L. (2003). Integrating the principles of energy conservation during everyday activities. *Caring*, 22(1), 30-31.
- Brollier, C., Hamrick, N., & Jacobson, B. (1993). Aerobic exercise: A potential occupational therapy modality for adolescents with depression. *Occupational Therapy in Mental Health*, 12(4), 19-29.
- Brown, C., Goetz, J., Van Sciver, A., Sullivan, D., & Hamera, E. (2006). A psychiatric rehabilitation approach to weight loss. *Psychiatric Rehabilitation Journal*, 29, 267-273. (Level II)
- Burr, N., Pratt, A. L., & Smith, P. J. (2002). An alternative splinting and rehabilitation protocol for metacarpophalangeal joint arthroplasty in patients with rheumatoid arthritis. *Journal of Hand Therapy*, 15(1), 41-47. (Level III)
- Burtner, P. A., Anderson, J. B., Marcum, M. L., Poole, J. L., Qualls, C., & Picchiarini, M. S. (2003). A comparison of static and dynamic wrist splints using electromyography in individuals with rheumatoid arthritis. *Journal of Hand Therapy*, 16, 320-325. (Level III)

- Burtner, P. A., Dukeminier, A., Ben, L., Qualls, C., & Scott, K. (2006). Visual perceptual skills and related school functions in children with hemiplegic cerebral palsy. *New Zealand Journal of Occupational Therapy*, 53(1), 24–29. (Level II)
- Callinan, N. J., & Mathiowetz, V. (1996). Soft versus hard resting hand splints in rheumatoid arthritis: Pain relief, preference, and compliance. *American Journal of Occupational Therapy*, 50(5), 347-353.
- Campbell, A. J., Robertson, M. C., La Grow, S. J., Kerse, N. M., Sunderson, G. F., Jacobs, R. J., . . . Hale, L. A. (2005). Randomised controlled trial of prevention of falls in people aged >75 with severe visual impairments: The VIP trial. *BMJ*, 331(7520), 817–824. (Level I)
- Carr, S. H. (1991). Adaptation to loss: Occupational therapy as a way of life. *American Journal of Occupational Therapy*, 45(2), 167-170.
- Carter, A. C. (1999). Occupational therapy perspective... breast cancer reconstructive rehabilitation. *Journal of Bodywork & Movement Therapies*, 3(1), 19-23.
- Cate, Y., & Richards, L. (2000). Relationship between performance on test of basic visual functions and visual-perceptual processing in persons after brain injury. *American Journal of Occupational Therapy*, 54, 326–334. (Level II)
- Cate, Y., Baker, S. S., & Gilbert, M. P. (1995). Occupational therapy and the person with diabetes and vision impairment. *American Journal of Occupational Therapy*, 49(9), 905-911.
- Chan, M. L. (1998). The role of occupational therapy in rheumatoid arthritis management. *Annals of the Academy of Medicine, Singapore*, 27(1), 120-121.
- Chan, S. (2004). Chronic obstructive pulmonary disease and engagement in occupation. *American Journal of Occupational Therapy*, 58(4), 408-415.
- Chard, G., Liu, L., & Mulholland, S. (2009). Verbal cueing and environmental modifications: Strategies to improve engagement in occupations in persons with Alzheimer disease. *Physical and Occupational Therapy in Geriatrics*, 27, 197–211. (Level III)
- Cheng, S. K. W., & Man, D. W. K. (2006). Management of impaired self-awareness in persons with traumatic brain injury. *Brain Injury*, 20, 621–628. (Level II)
- Cheville, A. (2001). Rehabilitation of patients with advanced cancer. *Cancer*, 92(4 SUPPL), 1039-1048.
- Cheville, A. L., Troxel, A. B., Basford, J. R., & Kornblith, A. B. (2008). Prevalence and treatment patterns of physical impairments in patients with metastatic breast cancer. *Journal of Clinical Oncology*, 26(16), 2621-2629.
- Chida, S., Shimada, Y., Matsunaga, T., Sato, M., Hatakeyama, K., & Mizoi, K. (2002). Occupational therapy for accessory nerve palsy after radical neck dissection. *Tohoku Journal of Experimental Medicine*, 196(3), 157-165.
- Chong, C. S. (2008). Management strategies for post-concussion syndrome after mild head injury: A systematic review. *Hong Kong Journal of Occupational Therapy*, 18(2), 59–67. (Systematic review)
- Cicerone, K. D., Mott, T., Azolay, J., & Friel, J. C. (2004). Community integration and satisfaction with functioning after intensive cognitive rehabilitation for traumatic brain injury. *Archives of Physical Medicine and Rehabilitation*, 86, 943– 950. (Level II)
- Clark, B. M. (2000). Rheumatology: 9. physical and occupational therapy in the management of arthritis. *CMAJ Canadian Medical Association Journal*, 163(8), 999-1005.
- Clark, F., Azen, S. P., Zemke, R., Jackson, J., Carlson, M., Mandel, D., ... Lipson, L. (1997). Occupational therapy for independent-living older adults. A randomized controlled trial. *JAMA*, 278(16), 1321–1326.
- Clark, F., Jackson, J., Carlson, M., Chou, C.-P., Cherry, B. J., Jordan-Marsh, M., ... Azen, S. P. (2012). Effectiveness of a lifestyle intervention in promoting the well-being of independently living older people: results of the Well Elderly 2

Randomised Controlled Trial. *Journal of Epidemiology and Community Health*, 66(9), 782–790.
<https://doi.org/10.1136/jech.2009.099754>

- Clarke, C. E., Furnston, A., Morgan, E., Patel, S., Sackley, C., Walker, M. F., . . . Wheatley, K. (2009). Pilot randomized controlled trial of occupational therapy to optimize independence in Parkinson's disease: The PD OT trial. *Journal of Neurology, Neurosurgery and Psychiatry*, 80, 976–978. (Level I)
- Clemson, L., Cumming, R. G., Kendig, H., Swann, M., Heard, R., & Taylor, K. (2004). The effectiveness of a community-based program for reducing the incidence of falls in the elderly: A randomized trial. *Journal of American Geriatrics Society*, 52, 1487–1494. (Level I)
- Cochran, A. (2007). Client-centered practice for agricultural workers with osteoarthritis. *OT Practice*, 12(12), 17-22.
- Cook, A., & Burkhardt, A. (1994). The effect of cancer diagnosis and treatment on hand function. *American Journal of Occupational Therapy*, 48(9), 836-839.
- Cooper, J. (1998). Occupational therapy intervention with radiation-induced brachial plexopathy. *European Journal of Cancer Care*, 7(2), 88-92.
- Cox, D. L. (2002). Chronic fatigue syndrome: An evaluation of an occupational therapy inpatient intervention. *British Journal of Occupational Therapy*, 65, 461–468. (Level II)
- Cranitch, C. (2003). Conducting a needs assessment to justify the provision of occupational therapy services in a rheumatology outpatient clinic. *Australian Occupational Therapy Journal*, 50(1), 23-29.
- Crennan, M., & MacRae, A. (2010). Occupational Therapy Discharge Assessment of Elderly Patients from Acute Care Hospitals. *Physical & Occupational Therapy In Geriatrics*, 28(1), 33–43. <https://doi.org/10.3109/02703180903381060>
- Crompton, S. (2004). *Occupational therapy intervention in cancer: Guidance for professionals, managers and decision-makers*. Southwark, London, United Kingdom: College of Occupational Therapists Ltd.
- Crowe, J., & Henderson, J. (2003). Pre-arthroplasty rehabilitation is effective in reducing hospital stay. *Canadian Journal of Occupational Therapy*, 70, 88–96. (Level I)
- Cumming, R. G., Thomas, M., Szonjy, G., Frampton, G., Salkeld, G., & Clemson, L. (2001). Adherence to occupational therapist recommendations for home modifications for falls prevention. *American Journal of Occupational Therapy*, 55, 641–648. (Level III)
- Curtin, C. (1991). Psychosocial intervention with an adolescent with diabetes using the model of human occupation. *Occupational Therapy in Mental Health*, 11(2-3), 23-36.
- Custer, V. L., & Wassink, K. E. (1991). Occupational therapy intervention for an adult with depression and suicidal tendencies. *American Journal of Occupational Therapy*, 45(9), 845-848.
- Dahlin-Ivanoff, S., Sonn, U., & Sveinsson, E. (2002). A health education program for elderly persons with visual impairments and perceived security in the performance of daily occupations: A randomized study. *American Journal of Occupational Therapy*, 56(3), 322–330. (Level I)
- Daus, C. (1998). Helping COPD patients help themselves... occupational therapy. *RT: The Journal for Respiratory Care Practitioners*, 11(2), 85-88.
- Davison, J., Bond, J., Dawson, P., Steen, I. M., & Kenny, R. A. (2005). Patients with recurrent falls attending Accident & Emergency benefit from multifactorial intervention: A randomized controlled trial. *Age and Ageing*, 34, 162–168. (Level I)
- Dawson, S., & Barker, J. (1995). Hospice and palliative care: A delphi survey of occupational therapists' roles and training needs. *Australian Occupational Therapy Journal*, 42(3), 119-127.

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- de Craen, A. J. M., Westendorp, R. G. J., Willems, C. G., Buskens, I. C. M., & Gusssekloo, J. (2006). Assistive devices and community-based services among 85-year-old community-dwelling elderly in The Netherlands: ownership, use, and need for intervention. *Disability and Rehabilitation. Assistive Technology*, 1(3), 199–203.
- deBuck, P. M., leCessie, S., vandenHout, W., Peeters, A., Ronday, H., Westedt, M., et al. (2005). Randomized comparison of a multidisciplinary job-retention vocational rehabilitation program with usual outpatient care in patients with chronic arthritis at risk for job loss. *Arthritis & Rheumatism: Arthritis Care & Research*, 53(5), 682-690.
- Dellhag, B., Hosseini, N., Bremell, T., & Ingvarsson, P. E. (2001). Disturbed grip function in women with rheumatoid arthritis. *Journal of Rheumatology*, 28, 2624–2633. (Level II)
- Derom, E., Marchand, E., & Troosters, T. (2007). Pulmonary rehabilitation in chronic obstructive pulmonary disease. *Annales De Readaptation Et De Medecine Physique*, 50(7), 615-626.
- Dettenmeier, P. A. (1990). Planning for successful home mechanical ventilation. *AACN Clinical Issues in Critical Care Nursing*, 1(2), 267-279.
- Devereaux, E., & Carlson, M. (1992). The role of occupational therapy in the management of depression. *American Journal of Occupational Therapy*, 46(2), 175-180.
- Dirette, D. (2004). A comparison of attention, processing and strategy use by adults with and without acquired brain injuries. *Brain Injury*, 18, 1219–1227. (Level II)
- Dohnke, B., Knäuper, B., & Müller-Fahrnow, W. (2005). Perceived self-efficacy gained from, and health effects of, a rehabilitation program after hip joint replacement. *Arthritis Rheum*, 53(4), 585-592.
- Dooley, N. R., & Hinojosa, J. (2004). Improving quality of life for persons with Alzheimer's disease and their family caregivers: Brief occupational therapy intervention. *American Journal of Occupational Therapy*, 58, 561–569. (Level III)
- Driessen, M., Dekker, J., Lankhorst, G., & van der Zee, J. (1997). Occupational therapy for patients with chronic diseases: CVA, rheumatoid arthritis and progressive diseases of the central nervous system. *Disability & Rehabilitation*, 19(5), 198204.
- Dubouloz, C., Laporte, D., Hall, M., Ashe, B., & Smith, C. D. (2004). Transformation of meaning perspectives in clients with rheumatoid arthritis. *American Journal of Occupational Therapy*, 58(4), 398-407.
- Dumoulin, C., Korner-Bitensky, N., & Tannenbaum, C. (2005). Urinary incontinence after stroke: Does rehabilitation make a difference? A systematic review of the effectiveness of behavioral therapy. *Topics in Stroke Rehabilitation*, 12(3), 66–76. (Systematic review)
- Eckloff, S. G., & Thornton, B. C. (2002). Prescribing assistive devices for patients with rheumatoid arthritis: Careful selection of equipment helps patients perform daily functions. *Journal of Musculoskeletal Medicine*, 19(1), 27-34.
- Egan, M. Y., & Brousseau, L. (2007). Splinting for osteoarthritis of the carpometacarpal joint: A review of the evidence. *American Journal of Occupational Therapy*, 61(1), 70-78.
- Egan, M., Brousseau, L., Farmer, M., Ouimet, M., Rees, S., Tugwell, P., et al. (2006). Splints and orthosis for treating rheumatoid arthritis. *Cochrane Database of Systematic Reviews*, 4
- Egan, M., Kessler, D., Laporte, L., Metcalfe, V., & Carter, M. (2007). A pilot randomized controlled trial of community-based occupational therapy in late stroke rehabilitation. *Topics in Stroke Rehabilitation*, 14(5), 37–45. (Level I)
- Eklund, K., Sjöstrand, J., & Dahlin-Ivanoff, S. (2008). A randomized controlled trial of a health-promotion programme and its effect on ADL dependence and self-reported health problems for the elderly visually impaired. *Scandinavian Journal of Occupational Therapy*, 15, 68–74. (Level I)
- Eklund, K., Sonn, U., & Dahlin-Ivanoff, S. (2004). Long-term evaluation of a health education programme for elderly persons with visual impairment: A randomized study. *Disability and Rehabilitation*, 26, 401–409. (Level I)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale:

Via Angelo Emo 144, 00136 Roma

Email / sito web:

info@aito.it / www.aito.it

- Elinge, E., Lofgren, B., Gagerman, E., & Nyberg, L. (2003). A group learning programme for older people with hip fractures: A randomized study. *Scandinavian Journal of Occupational Therapy*, 10(1), 27–33. (Level I)
- Estabrooks, P. A., Nelson, C. C., Xu, S., King, D., Bayliss, E. A., Gaglio, B., . . . Glasgow, R. E. (2005). The frequency and behavioral outcomes of goal choices in the self-management of diabetes. *Diabetes Educator*, 31, 391–400. (Level I)
- Evans, L., & Brewis, C. (2008). The efficacy of community-based rehabilitation programmes for adults with TBI. *International Journal of Therapy and Rehabilitation*, 15, 446–457. (Systematic review)
- Farina, E., Mantovani, F., Fioravanti, R., Rotella, G., Villanelli, F., Imborone, E., . . . Postoglione, A. (2006). Efficacy of recreational and occupational activities associated to psychologic support in mild to moderate Alzheimer disease: A multicenter controlled study. *Alzheimer Disease & Associated Disorders*, 20, 275–282. (Level II)
- Findley, L. J., & Cox, D. L. (2000). Severe and very severe patients with chronic fatigue syndrome: Perceived outcome following an inpatient programme. *Journal of Chronic Fatigue Syndrome*, 7(3), 33–47. (Level II)
- Fine, J. (2001). The effect of leisure activity on depression in the elderly: Implications for the field of occupational therapy. *Occupational Therapy in Health Care*, 13(1), 45–59.
- Finlayson, M. (2005). Pilot study on an energy conservation education program delivered by telephone conference call to people with multiple sclerosis. *NeuroRehabilitation*, 20, 267–277. (Level III)
- Finlayson, M. L., Peterson, E. W., & Cho, C. (2009). Pilot study of a fall risk management program for middle aged and older adults with MS. *NeuroRehabilitation*, 25, 107–115. (Level III)
- Finnerty, J. P., Keeping, I., Bullough, I., & Jones, J. (2001). The effectiveness of outpatient pulmonary rehabilitation in chronic lung disease: A randomized controlled trial. *Chest*, 119, 1705–1710. (Level I)
- Fong, K. N. K., & Howie, D. R. (2009). Effects of an explicit problem-solving skills training program using a metacomponential approach for outpatients with acquired brain injury. *American Journal of Occupational Therapy*, 63, 525–534. (Level II)
- Frederick, J. T., Steinman, L. E., Prohaska, T., Satariano, W. A., Bruce, M., Bryant, L., et al. (2007). Community-based treatment of late life depression. an expert panel informed literature review. *American Journal of Preventive Medicine*, 33(3), 222–249.
- Fuller, H. M., Thomas, J. J., & Rice, M. S. (2006). Perceived risk: Effects of reaching and placing performance in persons with cerebrovascular accident. *American Journal of Occupational Therapy*, 60, 379–387. (Level III)
- Funaki, Y., Kaneko, F., & Okamura, H. (2005). Study on factors associated with changes in quality of life of demented elderly persons in group homes. *Scandinavian Journal of Occupational Therapy*, 12(1), 4–9. (Level III)
- Garrett, K., & Kirchner, S. (1995). Oncologic emergencies -- the role of physical therapy and occupational therapy. *Rehabilitation Oncology*, 13(1), 10–24.
- Garvey, J., Connolly, D., Boland, F., & Smith, S. M. (2015). OPTIMAL, an occupational therapy led self-management support programme for people with multimorbidity in primary care: a randomized controlled trial. *BMC family practice*, 16(1), 1.
- Gasser-Wieland, T. L., & Rice, M. S. (2002). Occupational embeddedness during a reaching and placing task with survivors of cerebral vascular accident. *OTJR: Occupation, Participation and Health*, 22(4), 153–160. (Level I)
- Gier, M. D., Levick, M. D., & Blazina, P. J. (1988). Stress reduction with heart transplant patients and their families: A multidisciplinary approach. *Journal of Heart Transplantation*, 7(5), 342–347.
- Gil, N., & Josman, N. (2001). Memory and metamemory performance in Alzheimer's disease and healthy elderly: The Contextual Memory Test (CMT). *Aging and Clinical Experimental Research*, 13, 309–315. (Level II)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Gilbertson, L., & Langhorne, P. (2000). Home-based occupational therapy: Stroke patients' satisfaction with occupational performance and service provision. *British Journal of Occupational Therapy*, *63*, 464–468. (Level III)
- Gilbertson, L., Langhorne, P., & Walker, A. (2000). Domiciliary occupational therapy for patients with stroke discharged from hospital: Randomised controlled trial. *BMJ*, *320*, 603–606. (Level I)
- Gilbey, H., Ackland, T. R., Wang, A.W., Morton, A. R., Trouchet, T., & Tapper, J. (2003). Exercise improves early functional recovery after total hip arthroplasty. *Clinical Orthopaedics and Related Research*, *408*, 193–200. (Level I)
- Gillen, G., Berger, S., Lotia, S., Morreale, J., Siber, M., & Trudo, W. (2007). Improving community skills after lower extremity joint replacement. *Physical & Occupational Therapy in Geriatrics*, *25*(4), 41–54. (Level III)
- Gitlin, L. M., Hauck, W. W., Dennis, M. P., Winter, L., Hodgson, N., & Schinfeld, S. (2009). Long-term effect on mortality of a home intervention that reduces functional difficulties in older adults: Results from a randomized trial. *Journal of the American Geriatric Society*, *57*, 476–481. (Level I).
- Gitlin, L. N., Corcoran, M., Winter, L., Boyce, A., & Hauck, W. W. (2001). A randomized controlled trial of a home environmental intervention: Effect of efficacy and upset in caregivers and on daily function of persons with dementia. *Gerontologist*, *41*, 4–14. (Level I)
- Gitlin, L. N., Hauck, W. W., Dennis, M. P., & Winter, L. (2005). Maintenance of effects of the home environment skillbuilding program for family caregivers and individuals with Alzheimer's disease and related disorders. *The Journals of Gerontology*, *60*, 368–374. (Level I)
- Gitlin, L. N., Winter, L., Burke, J., Chernet, N., Dennis, M. P., & Hauck, W. W. (2008). Tailored activities to manage neuropsychiatric behaviors in persons with dementia and reduced caregiver burden: A randomized pilot study. *American Journal of Geriatric Psychiatry*, *16*, 229–239. (Level I)
- Gitlin, L. N., Winter, L., Dennis, M. P., & Hauck, W. W. (2007). A non-pharmacological intervention to manage behavioral and psychological symptoms of dementia and reduce caregiver distress: Design and methods of Project ACT. *Clinical Interventions in Aging*, *2*, 695–703. (Level I)
- Gitlin, L., Winter, L., Dennis, M., Corcoran, M., Schinfeld, S., & Hauck, W. (2006). A randomized trial of a multicomponent home intervention to reduce functional difficulties in older adults. *Journal of the American Geriatric Society*, *54*, 809–816. (Level I)
- Giuffrida, C. G., Demery, J. A., Reyes, L. R., Lebowitz, B. K., & Hanlon, R. E. (2009). Functional skill training in men with traumatic brain injury. *American Journal of Occupational Therapy*, *63*, 396–405. (Level II)
- Glasgow, R. E., Nutting, P. A., King, D. K., Nelson, C. C., Cutter, G., Gaglio, B., . . . Amthauer, H. (2004). A practical randomized trial to improve diabetes care. *Journal of General Internal Medicine*, *19*, 1167–1174. (Level I)
- Glasgow, R. E., Nutting, P. A., King, D. K., Nelson, C. C., Cutter, G., Gaglio, B., . . . Whitesides, H. (2005). Randomized effectiveness trial of a computer-assisted intervention to improve diabetes care. *Diabetes Care*, *28*(1), 33–39. (Level I)
- Glasgow, R. E., Nutting, P. A., Toobert, D. J., King, D. K., Strycker, L. A., Jex, M., . . . Merenich, J. (2006). Effects of a brief computer-assisted diabetes self-management intervention on dietary, biological and quality-of-life outcomes. *Chronic Illness*, *2*(1), 27–38. (Level I)
- Glasgow, R. E., Strycker, L. A., King, D. K., Toobert, D. J., Rahm, A. K., Jex, M., & Nutting, P. A. (2006). Robustness of a computer-assisted diabetes self-management intervention across patient characteristics, healthcare settings, and intervention staff. *American Journal of Managed Care*, *13*, 137–145. (Level I)
- Goldstein, N. E., Genden, E., & Morrison, R. S. (2008). Palliative care for patients with head and neck cancer: "I would like a quick return to a normal lifestyle". *JAMA - Journal of the American Medical Association*, *299*(15), 1818–1825.

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Gosman-Hedström, G., Claesson, L., & Blomstrand, C. (2002). Assistive devices in elderly people after stroke: A longitudinal, randomized study—The Göteborg 70+ stroke study. *Scandinavian Journal of Occupational Therapy*, 9, 109–118. (Level I)
- Goto, Y., Kurosawa, H., Mori, N., Kurokawa, Y., Hida, W., & Kohzuki, M. (2004). Improved activities of daily living, psychological state and health-related quality of life for 12 months following lung volume reduction surgery in patients with severe emphysema. *Respirology*, 9, 337–344. (Level II)
- Goverover, Y., & Hinojosa, J. (2004). Interrater reliability and discriminant validity of the Deductive Reasoning Test. *American Journal of Occupational Therapy*, 58(1), 104–108. (Level II)
- Graff, M. J. L., Adang, E. M. M., Vernooij-Dassen, M. J. M., Dekker, J., Jönsson, L., Thijssen, M., . . . OldeRikkert, M. G. M. (2008). Community occupational therapy for older patients with dementia and their care givers: Cost effectiveness study. *BMJ*, 336(7636) 134–138. (Level III)
- Graff, M. J. L., Vernooij-Dassen, M. J. F. J., Hoefnagel, W. H. L., Dekker, J., & de Witte, L. P. (2003). Occupational therapy at home for older individuals with mild to moderate cognitive impairments and their primary caregivers: A pilot study. *OTJR: Occupation, Participation and Health*, 23, 155–164. (Level III)
- Graff, M. J. L., Vernooij-Dassen, M. J. M., Thijssen, M., Dekker, J., Hoefnagels, W. H. L., & OldeRikkert, M. G. M. (2007). Effects of community occupational therapy on quality of life, mood, and health status in dementia patients and their caregivers: A randomized controlled trial. *Journals of Gerontology A: Biological Sciences and Medical Sciences*, 62, 1002–1009. (Level I)
- Grant, M. (2001). Mothers with arthritis, child care and occupational therapy: Insight through case studies. *British Journal of Occupational Therapy*, 64(7), 322-329.
- Grant, M. (2005). Occupational therapy for people with osteoarthritis: Scope of practice and evidence base... including commentary by hammond A and rogers JC. *International Journal of Therapy & Rehabilitation*, 12(1), 7-13.
- Griffiths, T. L., Burr, M. L., Campbell, I. A., LewisJenkins, V., Mullins, J., Shiels, K., et al. (2000). Results at 1 year of outpatient multidisciplinary pulmonary rehabilitation: A randomised controlled trial. *Lancet*, 355(9201), 362-368.
- Gunnarsson, A. B., Jansson, J. -, & Eklund, M. (2006). The tree theme method in psychosocial occupational therapy: A case study. *Scandinavian Journal of Occupational Therapy*, 13(4), 229-240.
- Haber, D., Looney, C., Babola, K., Hinman, M., & Utsey, C. J. (2000). Impact of a health promotion course on inactive, overweight, or physically limited older adults. *Family and Community Health*, 22(4), 48–56. (Level III)
- Hadjistavropoulos, H., Asmundson, G. J. G., LaChapelle, D. L., & Quine, A. (2002). The role of health anxiety among patients with chronic pain in determining response to therapy. *Pain Research & Management*, 7, 127–133. (Level III)
- Hagsten, B., Svensson, O., & Gardulf, A. (2004). Early individualized postoperative occupational therapy training in 100 patients improves ADL after hip fracture: A randomized trial. *Acta Orthopaedica Scandinavica*, 75, 177–183. (Level I)
- Hagsten, B., Svensson, O., & Gardulf, A. (2006). Health-related quality of life and self-reported ability concerning ADL and IADL after hip fracture: A randomized trial. *Acta Orthopaedica*, 77(1), 714–719. (Level I)
- Haig, A. J., Geisser, M. E., Nicholson, C., Parker, E., Yoamkawa, K., Montgomery, D., & Booker, E. (2003). The effect of order on testing in functional performance in persons with and without chronic back pain. *Journal of Occupational Rehabilitation*, 13, 115–123. (Level III)
- Hale, S., Michalak, E. E., Hayashi, B., & Lam, R. W. (2005). Relief of chronic or resistant depression (re-ChORD): A clinical research program for chronic depression. *Occupational Therapy Now*, 7(3), 28-30.
- Hällgren, M., & Kottorp, A. (2005). Effects of occupational therapy intervention in activities of daily living and awareness of disability in persons with intellectual disabilities. *Australian Occupational Therapy Journal*, 52, 350–359. (Level III)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Hammell, K. W. (2015). Quality of life, participation and occupational rights: A capabilities perspective. *Australian Occupational Therapy Journal*, 62(2), 78–85. <https://doi.org/10.1111/1440-1630.12183>
- Hammond, A. (1994). Joint protection behavior in patients with rheumatoid arthritis following an education program. *Arthritis Care and Research*, 7(1), 5-9.
- Hammond, A. (2004). What is the role of the occupational therapist?. *Best Practice and Research in Clinical Rheumatology*, 18(4), 491-505.
- Hammond, A. (2008). Rehabilitation in musculoskeletal diseases. *Best Practice and Research: Clinical Rheumatology*, 22(3), 435-449.
- Hammond, A., & Freeman, K. (2001). One-year outcomes of a randomized controlled trial of an educational-behavioural joint protection programme for people with rheumatoid arthritis. *Rheumatology (Oxford, England)*, 40(9), 1044-1051.
- Hammond, A., & Freeman, K. (2004). The long-term outcomes from a randomized controlled trial of an educational-behavioural joint protection programme for people with rheumatoid arthritis. *Clinical Rehabilitation*, 18(5), 520-528.
- Hammond, A., & Freeman, K. (2006). Community patient education and exercise for people with fibromyalgia: A parallel group randomized controlled trial. *Clinical Rehabilitation*, 20, 835–846. (Level I)
- Hammond, A., Jeffreson, P., Jones, N., Gallagher, J., & Jones, T. (2002). Clinical applicability of an educational-behavioural joint protection programme for people with rheumatoid arthritis. *British Journal of Occupational Therapy*, 65(9), 405-412.
- Hammond, A., Young, A., & Kidao, R. (2004). A randomised controlled trial of occupational therapy for people with early rheumatoid arthritis. *Annals of the Rheumatic Diseases*, 63(1), 23-30.
- Harris, S. R., & Mortenson, P. A. (2006). Playfulness in children with traumatic brain injury: A preliminary study. *Physical & Occupational Therapy in Pediatrics*, 26, 181–198. (Level II)
- Harrison-Paul, J., & Drummond, A. E. R. (2006). A randomized controlled trial of occupational therapy in oncology: Challenges in conducting a pilot study. *British Journal of Occupational Therapy*, 69, 130–133. (Level I)
- Hartman-Maeir, A., Soroker, N., Ring, H., & Katz, N. (2002). Awareness of deficits in stroke rehabilitation. *Journal of Rehabilitation Medicine*, 34, 158–164. (Level III)
- Haskett, S., Backman, C., Porter, B., Goyert, J., & Palejko, G. (2004). A crossover trial of custom-made and commercially available wrist splints in adults with inflammatory arthritis. *Arthritis Care and Research*, 51(5), 792-799.
- Hatchard, K., & Missiuna, C. (2003). An occupational therapist's journey through bipolar affective disorder. *Occupational Therapy in Mental Health*, 19(2), 1-17.
- Hay, J., LaBree, L., Luo, R., Clark, F., Carlson, M., Mandel, D., ...Azen, S. P. (2002). Cost-effectiveness of preventive occupational therapy for independent-living older adults. *Journal of the American Geriatrics Society*, 50(8), 1381–1388.
- Heather, F. (2002). Pro-motion: A positive way forward for clients with severe and enduring mental health problems living in the community, part 1. *British Journal of Occupational Therapy*, 65(12), 551-558.
- Heaton, J., McMurray, R., Sloper, P., & Nettleton, S. (2000). Rehabilitation and total hip replacement: Patients' perspectives on provision. *International Journal of Rehabilitation Research*, 23(4), 253-259.
- Helewa, A., Goldsmith, C. H., Lee, P., Bombardier, C., Hanes, B., Smythe, H. A., et al. (1991). Effects of occupational therapy home service on patients with rheumatoid arthritis. *Lancet*, 337(8755), 1453-1456.
- Hendriks, M. R., Evers, S. M., Bleijlevens, M. H., Van Haastregt, J. C., Crebolder, H. F., & Van Eijk, J. T. (2008). Cost-effectiveness of a multidisciplinary fall prevention program in community-dwelling elderly people: A randomized controlled trial (ISRCTN64716113). *International Journal of Technology Assessment in Health Care*, 24, 193–202. (Level I)

- Hendriksen, H., & Harrison, R. A. (2001). Occupational therapy in accident and emergency departments: A randomized controlled trial. *Journal of Advanced Nursing*, 36, 727–732. (Level I)
- Horne, D., & Corsello, P. (1993). Physical and occupational therapy for patients with chronic lung disease. *Seminars in Respiratory Medicine*, 14(6), 466–481.
- Huang, M. E., Wartella, J. E., & Kreuzer, J. S. (2001). Functional outcomes and quality of life in patients with brain tumors: A preliminary report. *Archives of Physical Medicine and Rehabilitation*, 82, 1540–1546. (Level III)
- Imms, C. (2004). Occupational performance challenges for children with congenital heart disease: A literature review. *Canadian Journal of Occupational Therapy*, 71(3), 161–172.
- Ip, S. P. S., Leung, Y. F., & Choy, K. L. (2004). Short-stay in-patient rehabilitation of elderly patients with chronic obstructive pulmonary disease: Prospective study. *Hong Kong Medical Journal*, 10, 312–318. (Level II)
- Iyengar, K. P., Nadkarni, J. B., Ivanovic, N., & Mahale, A. (2007). Targeted early rehabilitation at home after total hip and knee joint replacement: Does it work? *Disability and Rehabilitation*, 29, 495–502. (Level III)
- Jackson, J. P., & Schkade, J. K. (2001). Occupational adaptation model versus biomechanical-rehabilitation model in the treatment of patients with hip fractures. *American Journal of Occupational Therapy*, 55, 531–537. (Level II)
- Jackson, N. C. (1991). Pulmonary rehabilitation for mechanically ventilated patients. *Critical Care Nursing Clinics of North America*, 3(4), 591–600.
- Johansson, A., & Björklund, A. (2016). The impact of occupational therapy and lifestyle interventions on older persons' health, well-being, and occupational adaptation: A mixed-design study. *Scandinavian Journal of Occupational Therapy*, 23(3), 207–219. <https://doi.org/10.3109/11038128.2015.1093544>
- Johnson, N. J., & Grindler, D. (1990). Home rehabilitation for chronic obstructive pulmonary disease: A physical and occupational therapy approach. *Journal of Home Health Care Practice*, 2(2), 19–43.
- Jones, F. A., Rodger, S., Ziviani, J., & Boyd, R. (2009). Transition to secondary school for students with cerebral palsy and implications for quality of life: A systematic review. *Journal of Occupational Therapy Schools & Early Intervention*, 2, 133–147. (Systematic review)
- Jongbloed, L., & Wendland, T. (2002). The impact of reimbursement systems on occupational therapy practice in Canada and the United States of America. *Canadian Journal of Occupational Therapy. Revue Canadienne D'ergotherapie*, 69(3), 143–152.
- Kamble, R., Goswami, S., Saigal, J., Sarin, R., & Jalali, R. (2003). Prospective neuro-cognitive assessment (using LOTCA) and quality of life (QOL) and activities of daily living (ADL) in children with brain tumour treated with surgery and localized high-precision radiotherapy. *Indian Journal of Occupational Therapy*, 35(3), 10–12. (Level III)
- Kealey, P., & McIntyre, I. (2005). An evaluation of the domiciliary occupational therapy service in palliative cancer care in a community trust: A patient and carers perspective. *European Journal of Cancer Care*, 14(3), 232–243.
- Kendig, T. D., Nonailhada, J., Ramaker, J., & Samuels, F. (2008). Management of a patient with lung metastases requiring prolonged mechanical ventilation (PMV):
- Kielhofner, G., Braveman, B., Fogg, L., & Levin, M. (2008). A controlled study of services to enhance productive participation among people with HIV/AIDS. *American Journal of Occupational Therapy*, 62(1), 36–54. (Level II)
- King, D. K., Estabrooks, P. A., Strycker, L. A., Toobert, D. J., Bull, S. S., & Glasgow, R. E. (2006). Outcomes of a multifaceted physical activity regimen as part of a diabetes self-management intervention. *Annals of Behavioral Medicine*, 31, 128–137. (Level I)
- Kos, D., Duportail, M., D'hooghe, M., Nagels, G., & Kerckhofs, E. (2007). Multidisciplinary fatigue management programme in multiple sclerosis: A randomized clinical trial. *Multiple Sclerosis*, 13, 996–1003. (Level I)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Kraaimaat, F. W., Brons, M. R., Geenen, R., & Bijlsma, J. W. (1995). The effect of cognitive behavior therapy in patients with rheumatoid arthritis. *Behaviour Research & Therapy*, 33(5), 487-495.
- Kusznir, A., Scott, E., Cooke, R. G., & Young, L. T. (1996). Functional consequences of bipolar affective disorder: An occupational therapy perspective. *Canadian Journal of Occupational Therapy*, 63(5), 313-322.
- laCour, K., Josephsson, S., Tishelman, C., & Nygard, L. (2007). Experiences of engagement in creative activity at a palliative care facility. *Palliative & Supportive Care*, 5(3), 241-250.
- La Grow, S. J., Robertson, M. C., Campbell, A. J., Clarke, G. A., & Kerse, N. M. (2006). Reducing hazard related falls in people 75 years and older with significant visual impairment: How did a successful program work? *Injury Prevention*, 12, 296-301. (Level III)
- Labell, K. L., & Mihailidis, A. (2006). The use of automated prompting to facilitate handwashing in persons with dementia. *American Journal of Occupational Therapy*, 60, 442-450. (Level III)
- Lala, A. P., & Kinsella, E. A. (2011). A phenomenological inquiry into the embodied nature of occupation at end of life. *Canadian Journal of Occupational Therapy*, 78(4), 246-254.
- Lambeck, L. C., van Mechelen, W., Buijs, P. C., Loisel, P., & Anema, J. R. (2009). An integrated care program to prevent work disability due to chronic low back pain: A process evaluation within a randomized controlled trial. *BMC Musculoskeletal Disorders*, 10(147). doi:10.1186/1471-2474-10-147. (Level I)
- Larson, K. B. (1990). Activity patterns and life changes in people with depression. *American Journal of Occupational Therapy*, 44(10), 902-906.
- Lau, O. W. Y., Leung, L. N. Y., & Wang, L. O. L. (2002). Cognitive behavioural techniques for changing the coping skills of patients with chronic pain. *Hong Kong Journal of Occupational Therapy*, 12, 13-20. (Level III)
- Law, M., Steinwender, S., & Leclair, L. (1998). Occupation, Health and Well-Being. *Canadian Journal of Occupational Therapy*, 65(2), 81-91. <https://doi.org/10.1177/000841749806500204>
- Lee, W. T., Chan, H. F., & Wong, E. (2005). Improvement of feeding independence in end-stage cancer patients under palliative care--a prospective, uncontrolled study. *Supportive Care in Cancer*, 13(12), 1051-1056.
- Legg, L., & Langhorne, P. (2004). Rehabilitation therapy services for stroke patients living at home: Systematic review of randomized trials. *Lancet*, 363(9406), 352-356. (Systematic review)
- Leland, N. E., Crum, K., Phipps, S., Roberts, P., & Gage, B. (2015). Advancing the value and quality of occupational therapy in health service delivery. *The American Journal of Occupational Therapy: Official Publication of the American Occupational Therapy Association*, 69(1), 6901090010p1-7. <https://doi.org/10.5014/ajot.2015.691001>
- Leslie, C. A. (2005). In the clinic. facing forward: How mental health occupational therapy offers a doorway to the world... for clients without a face. *OT Practice*, 10(11), 18-20.
- Leung, E. K. H. (2002). Evidence-based practice in occupational therapy. *Hong Kong Journal of Occupational Therapy*, 12, 21-32. (Systematic review)
- Li, L. C., & Iversen, M. D. (2005). Outcomes of patients with rheumatoid arthritis receiving rehabilitation. *Current Opinion in Rheumatology*, 17(2), 172-176. Li, L. C., Maetzel, A., Davis, A. M., Lineker, S. C., Bombardier, C., & Coyte, P. C. (2006). Primary therapist model for patients referred for rheumatoid arthritis rehabilitation: A cost-effectiveness analysis. *Arthritis Care and Research*, 55(3), 402-410.
- Li, L. C., Davis, A. M., Lineker, S. C., Coyte, P. C., & Bombardier, C. (2006). Effectiveness of the primary therapist model for rheumatoid arthritis rehabilitation: A randomized controlled trial. *Arthritis & Rheumatism*, 55(1), 4252.

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale:

Via Angelo Emo 144, 00136 Roma

Email / sito web:

info@aito.it / www.aito.it

- Li, L. C., Davis, A. M., Lineker, S., Coyte, P. C., & Bombardier, C. (2005). Outcomes of home-based rehabilitation provided by primary therapists for patients with rheumatoid arthritis: Pilot study. *Physiotherapy Canada*, 57(4), 255-264.
- Lieber, S. J., Rudy, T. E., & Boston, J. R. (2000). Effects of body mechanics training on performance of repetitive lifting. *American Journal of Occupational Therapy*, 54(7), 166-175. (Level III)
- Lindén, A., Boschian, K., Eker, C., Schalén, W., & Nordström, C.-H. (2004). Assessment of motor and process skills reflects brain-injured patients' ability to resume independent living better than neuropsychological tests. *Acta Neurologica Scandinavica*, 111, 48-53. (Level III)
- Linroth, R., Zander, S., Forde, S., Hanley, M., & Lins, J. (1996). Ramsey county day treatment services: Day treatment to extended day treatment centers to focus groups. *Occupational Therapy in Health Care*, 10(2), 89-99.
- Li-Tsang, C. W. P., Hung, L. K., & Mak, A. F. T. (2002). The effect of corrective splinting on flexion contracture of rheumatoid fingers. *Journal of Hand Therapy*, 15, 185-191. (Level II)
- Lloyd, C., & Coggles, L. (1988). Contribution of occupational therapy to pain management in cancer patient with metastatic breast disease. *American Journal of Hospice Care*, 5(6), 36-38.
- Lloyd, C., & Coggles, L. (1989). Occupational therapy and the cancer patient: An occupational behavior perspective. *Physical & Occupational Therapy in Geriatrics*, 7(4), 59-69.
- Lloyd, C., & Coggles, L. (1990). Psychosocial issues for people with cancer and their families. *Canadian Journal of Occupational Therapy*, 57(4), 211-215.
- Lloyd, C., & Neven, J. (1988). Occupational therapy and the elderly cancer patient. *Physical & Occupational Therapy in Geriatrics*, 6(3), 87-97.
- Logan, P. A., Coupland, C. A. C., Gladman, J. R. F., Sahota, O., Stoner-Hobbs, V., Robertson, K., ... Avery, A. J. (2010). Community falls prevention for people who call an emergency ambulance after a fall: randomised controlled trial. *BMJ*, 340(may11 1), c2102-c2102. <https://doi.org/10.1136/bmj.c2102>
- Logan, P. A., Gladman, J. R. F., Drummond, A. E. R., & Radford, K. A. (2003). A study of interventions and related outcomes in a randomized controlled trial of occupational therapy and leisure therapy for community stroke patients. *Clinical Rehabilitation*, 17, 249-255. (Level I)
- Lorenzi, C. M., Cilone, C., Rizzardi, R., Furino, V., Bellantone, T., Lugli, D., & Clini, E. (2004). Occupational therapy and pulmonary rehabilitation of disabled COPD patients. *Respiration*, 71, 246-251. (Level II)
- MacDonald, D. (2006). Occupational therapists: An environmental scan of the economic literature. *Ottawa, ON: CAOT Publications ACE. CIT0038*.
- MacDonald, R. (2004). Occupational therapy in rheumatology. *Therapy Weekly*, 30(27), 10-13.
- Malcus-Johnson, P., Carlqvist, C., Stureson, A., & Eberhardt, K. (2005). Occupational therapy during the first 10 years of rheumatoid arthritis. *Scandinavian Journal of Occupational Therapy*, 12(3), 128-135.
- Månsson, E., & Lexell, J. (2004) Performance of activities of daily living in multiple sclerosis. *Disability and Rehabilitation*, 26, 576-585. (Level III)
- Maravic, M., Bozonnat, M., Sevezan, A., Gasqueres, D., Pastor, J., Pere, M., et al. (2000). Preliminary evaluation of medical outcomes (including quality of life) and costs in incident RA cases receiving hospital-based multidisciplinary management. *Joint Bone Spine*, 67(5), 425-433.
- Mårtensson, L., Marklund, B., Baigi, A., Gunnarsson, M., & Fridlund, B. (2004). Long-term influences of a biopsychosocial rehabilitation programme for chronic pain patients. *Musculoskeletal Care*, 2, 152-164. (Level III)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Masiero, S., Boniolo, A., Wassermann, L., Machiedo, H., Volante, D., & Punzi, L. (2007). Effects of an educational-behavioral joint protection program on people with moderate to severe rheumatoid arthritis: A randomized controlled trial. *Clin Rheumatol*, 26(12), 2043-2050.
- Mastenbroek, I., & McGovern, L. (1991). The effectiveness of relaxation techniques in controlling chemotherapy induced nausea: A literature review. *Australian Occupational Therapy Journal*, 38(3), 137-142.
- Mathieux, R., Marotte, H., Battistini, L., Sarrazin, A., Bethier, M., & Masec, P. (2008). Early occupational therapy programme increases hand grip strength at 3 months: Results from a randomized, blind, controlled study in early rheumatoid arthritis. *Annals of the Rheumatic Diseases*, 68, 400-403. (Level I)
- Mathiowetz, V. G., Finlayson, M. L., Matuska, M., Chen, H. Y., & Luo, P. (2005). Randomized controlled trial of an energy conservation course for persons with multiple sclerosis. *Multiple Sclerosis*, 11, 592-601. (Level I)
- Mathiowetz, V. G., Matuska, K. M., Finlayson, M. L., Luo, P., & Chen, H. Y. (2007). One-year follow-up to a randomized controlled trial of an energy conservation course for persons with multiple sclerosis. *International Journal of Rehabilitation Research*, 30, 305-313. (Level I)
- Mathiowetz, V., Matuska, K. M., & Murphy, M. E. (2001). Efficacy of an energy conservation course for persons with multiple sclerosis. *Archives of Physical Medicine and Rehabilitation*, 82, 449-456. (Level I)
- McCulloch, J. (2007). Hospital anxiety and depression in myocardial infarction patients. *British Journal of Cardiology*, 14(2), 106-108.
- McDonald, S., Hetrick, S., & Green, S. (2008). Pre-operative education for hip or knee replacement. *Cochrane Database of Systematic Reviews*, 4
- McKay, D. A. (2002). Diabetes: Functional implications, general precautions, and practical considerations. *OT Practice*, 7(19), 11-17.
- McKay, H. G., King, D., Eakin, E. G., Seeley, J. R., & Glasgow, R. E. (2001). The diabetes network internet-based physical activity intervention: A randomized pilot study. *Diabetes Care*, 24, 1328-1334. (Level I)
- McLeod, B., & Mair, M. (2009). *Evaluation of city of Edinburgh council home care re-ablement service*. Scottish Government Social Research.
- McWha, J. L., Pachana, N. A., & Alpass, F. (2003). Exploring the therapeutic environment for older women with late-life depression: An examination of the benefits of an activity group for older people suffering from depression. *Australian Occupational Therapy Journal*, 50, 158-169. (Level III)
- Mead, K., Theadom, A., Byron, K., & Dupont, S. (2007). Pilot study of a 4-week pain coping strategies (PCS) programme for the chronic pain patient. *Disability and Rehabilitation*, 29, 199-203. (Level III)
- Mejer, E. M., Sluiter, J. K., Heyma, A., Sadiraj, K., & Frings-Dresen, M. H. W. (2006). Cost-effectiveness of multidisciplinary treatment in sick-listed patients with upper extremity musculoskeletal disorders: A randomized, controlled trial with one-year follow-up. *International Archives of Occupational & Environmental Health*, 79(8), 654-664. (Level I)
- Migliore, A. (2004). Improving dyspnea management in three adults with chronic obstructive pulmonary disease. *American Journal of Occupational Therapy*, 58(6), 639-646.
- Montgomery, P., & Bratton, G. N. (1992). The effect of an inpatient arthritis rehabilitation program on self-assessed functional ability. *Rehabilitation Nursing*, 17(6), 306-310.
- Moran, M. E. (2001). Osteoarthritis and occupational therapy intervention. *Physical Medicine & Rehabilitation*, 15(1), 65-81.
- Morgan, S. B., Kelkar, R., & Vyas, O. A. (2002). Client-centered occupational therapy for acute stroke patients. *Indian Journal of Occupational Therapy*, 34(1), 7-12. (Level I)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Morris, M. E., Perry, A., Bilney, B., Curran, A., Dodd, K., Wittwer, J. E., & Dalton, G. W. (2006). Outcomes of physical therapy, speech pathology, and occupational therapy for people with motor neuron disease: A systematic review. *Neurorehabilitation and Neural Repair*, 20, 424–434. (Systematic review)
- Mortenson, P. A., & Harris, S. R. (2006). Playfulness in children with traumatic brain injury: A preliminary study. *Physical & Occupational Therapy in Pediatrics*, 26, 181–198. (Level II)
- Mountain, G., & Chatters, R. (2016). The «Lifestyle Matters» Study: Results From a Trial of an Occupational Therapy Lifestyle Intervention for Older Adults. *American Journal of Occupational Therapy*, 70(4_Supplement_1), 7011515242p1. <https://doi.org/10.5014/ajot.2016.70S1-RP304A>
- Mozley, C. G. (2001). Exploring connections between occupation and mental health in care homes for older people. *Journal of Occupational Science*, 8(3), 14-19.
- Mozley, C. G., Schneider, J., Cordingley, L., Molineuz, M., Suggan, S., Hart, C., . . . Cruickshank, A. (2007). The Care Home Activity Project: Does introducing an occupational therapy programme reduce depression in care homes? *Aging & Mental Health*, 11(1), 99–107. (Level III)
- Nawate, Y., Kaneko, F., Hanaoka, H., & Okamura, H. (2008). Efficacy of group reminiscence therapy for elderly dementia patients residing at home: A preliminary report. *Physical & Occupational Therapy in Geriatrics*, 26(3), 57–68. (Level III)
- Neely, A. S., Vikström, S., & Josephsson, S. (2009). Collaborative memory intervention in dementia: Caregiver participation matters. *Neuropsychological Rehabilitation*, 29, 696–715. (Level I)
- Nesbit, S. G. (2004). My breast cancer: An occupational therapist's perspective. *Occupational Therapy in Mental Health*, 20(2), 51-67.
- Nieuwenhuisen, K., Bultmann, U., NeumeyerGromen, A., Verhoeven, A. C., Verbeek, J. H. A., & van der FeltzCornelis, C. M. (2008). Interventions to improve occupational health in depressed people. *Cochrane Database Syst.Rev.*, (2)
- Nilsson, I., Fitinghoff, H., & Lilja, M. (2007). Continuing to work after the onset of rheumatoid arthritis. *Work*, 28(4), 335-342.
- Nobili, A., Riva, E., Tettamanti, M., Lucca, U., Liscio, M., Petrucci, B., & Porro, G. S. (2004). The effect of a structured intervention on caregivers of patients with dementia and problem behaviors: A randomized controlled pilot study. *Alzheimer Disease and Associated Disorders*, 18, 75–82. (Level I)
- Nordmark, B., Blomqvist, P., Andersson, B., Hagerstrom, M., Nordh-Grate, K., Ronnqvist, R., et al. (2006). A two-year follow-up of work capacity in early rheumatoid arthritis: A study of multidisciplinary team care with emphasis on vocational support. *Scandinavian Journal of Rheumatology*, 35(1), 7-14.
- Norweg, A. M., Whiteson, J., Malgady, R., Mola, A., & Rey, M. (2005). The effectiveness of different combinations of pulmonary rehabilitation program components: A randomized controlled trial. *Chest*, 128, 663–672. (Level I)
- Norweg, A., Bose, P., Snow, G., & Berkowitz, M. E. (2008). A pilot study of a pulmonary rehabilitation programme evaluated by four adults with chronic obstructive pulmonary disease. *Occupational Therapy International*, 15(2), 114-132.
- Nyman, A. (2007). Influences of the social environment on engagement in occupations: The experience of persons with rheumatoid arthritis. *Scandinavian Journal of Occupational Therapy*, 14(1), 63.
- Oakley, F., Duran, L., Fisher, A., & Merritt, B. (2003). Differences in activities of daily living motor skills of persons with and without Alzheimer's disease. *Australian Occupational Therapy Journal*, 50(2), 72–78. (Level II)
- Oakley, F., Khin, N. A., Parks, R., Bauer, L., & Sanderland, T. (2002). Improvement in activities of daily living in elderly following treatment for post-bereavement depression. *Acta Psychiatrica Scandinavica*, 105, 231–234. (Level II)

- Oerlemans, H. M., Oostendorp, R. A. B., de Boo, T., Van der Laan, L., Severens, J. L., & Goris, R. H. A. (2000). Adjuvant physical therapy versus occupational therapy in patients with reflex sympathetic dystrophy/complex regional pain syndrome type 1. *Archives of Physical Medicine & Rehabilitation*, 81(1), 49–56. (Level II)
- Oertli, S. (2007). Cancer: Many diagnoses, one occupation-based approach. *OT Practice*, 12(20), 16-19.
- Öhman, A., Nygård, L., & Borell, L. (2001). The vocational situation in cases of memory deficits or younger-onset dementia. *Scandinavian Journal of Caring Sciences*, 15, 34–43. (Level III)
- Onor, M., Trevisiol, M., Negro, C., Alessandra, S., Saina, M., & Aguglia, E. (2007). Impact of a multimodal rehabilitative intervention on demented patients and their caregivers. *American Journal of Alzheimer's Disease & Other Dementias*, 22(4), 261–272. (Level III)
- Pagnotta, A., Baron, M., & Korner-Bitensky, N. (1998). The effect of a static wrist orthosis on hand function in individuals with rheumatoid arthritis. *The Journal of Rheumatology*, 25(5), 879-885.
- Pagnotta, A., Korner-Bitensky, N., Mazer, B., Baron, M., & Wood-Dauphinee, S. (2005). Static wrist splint use in the performance of daily activities by individuals with rheumatoid arthritis. *Journal of Rheumatology*, 32, 2136–2143. (Level II)
- Pai, V. B. (2000). Occupational therapy & its role in the rehabilitation of coronary heart disease (C.H.D.). *Asian Journal of Cardiovascular Nursing*, 8(2), 11-13.
- Palmer, P., & Simons, J. (1991). Joint protection: A critical review. *British Journal of Occupational Therapy*, 54(12), 453-458.
- Pardessus, V., Puisieux, F., Di Pompoe, C., Gaudefroy, C., Thevenon, A., & Dewailly, P. (2002). Benefits of home visits for falls and autonomy in the elderly: A randomized trial study. *American Journal of Physical Medicine & Rehabilitation*, 81(4), 247–252. (Level I)
- Pasquetto, E. (1995). Occupational therapy after the cure in a child with leukemia. *WFOT Bulletin*, 31, 19-21.
- PaternostroSluga, T., & Stieger, M. (2004). Hand splints in rehabilitation. *Critical Reviews in Physical and Rehabilitation Medicine*, 16(4), 233-256.
- Penfold, S. L. (1996). The role of the occupational therapist in oncology. *Cancer Treatment Reviews*, 22(1), 75-81.
- Persson, E., Rivano-Fischer, M., & Eklund, M. (2004). Evaluation of changes in occupational performance among patients in a pain management program. *Journal of Rehabilitation Medicine*, 36, 85–91. (Level III)
- Pigg, J. S. (1995). Rheumatoid arthritis: How allied health professionals can help... seventh in a special series of articles on diagnosis and management of rheumatoid arthritis. *Journal of Musculoskeletal Medicine*, 12(2), 27.
- Poole, J. L. (2000). A comparison of limb praxis abilities of persons with developmental dyspraxis and adult onset apraxia. *Occupational Therapy Journal of Research*, 20(7), 106–120. (Level II)
- Poole, J. L., Sadek, J., & Haaland, K. Y. (2009). Ipsilateral deficits in 1-handed shoe tying after left or right hemisphere stroke. *Archives of Physical Medicine and Rehabilitation*, 90, 1800–1805. (Level II)
- Pratt, A. L., & Burr, N. (2007). Post-operative rehabilitation after pip joint with early active motion: A retrospective review of outcomes. *British Journal of Hand Therapy*, 12(1), 22-27.
- Prkachin, K. M., Solomon, P., Hwang, T., & Mercer, S. R. (2001). Does experience influence judgements of pain behaviour? Evidence from relatives of pain patients and therapists. *Pain Research & Management*, 6, 105–112. (Level II)
- Query, M., & Taylor, R. R. (2005). Linkages between goal attainment and quality of life for individuals with chronic fatigue syndrome. *Occupational Therapy in Health Care*, 19(4), 3–22. (Level I)

- Rapoliene, J., & Krisciunas, A. (2006). The effectiveness of occupational therapy in restoring the functional state of hands in rheumatoid arthritis patients. *Medicina (Kaunas, Lithuania)*, *42*, 823–828. (Level II)
- Rashbaum, I., & Whyte, N. (1996). Occupational therapy in pulmonary rehabilitation: Energy conservation and work simplification techniques. *Physical Medicine & Rehabilitation Clinics of North America*, *7*(2), 325–340.
- Rathnayake, T. (2009). *Parkinson's disease: Occupational therapy-related treatment*. Adelaide, Australia: The Joanna Briggs Institute. (Systematic review)
- Reistetter, T. A., & Abreu, B. C. (2005). Appraising evidence on community integration following brain injury: A systematic review. *Occupational Therapy International*, *12*, 196–217. (Systematic review)
- Rice, M. S., Fertig, P., Maitra, K., & Miller, B. K. (2008). Reduced feedback: Motor learning strategy in persons with Alzheimer's disease. *Physical and Occupational Therapy in Geriatrics*, *27*, 122–138. (Level II)
- Rivard, A., Warren, S., Voaklander, D., & Jones, A. (2003). The efficacy of pre-operative home visits for total hip replacement clients. *Canadian Journal of Occupational Therapy*, *70*(4), 226–232.
- Roberts, P. S., & Robinson, M. R. (2014). Occupational therapy's role in preventing acute readmissions. *The American Journal of Occupational Therapy: Official Publication of the American Occupational Therapy Association*, *68*(3), 254–259. <https://doi.org/10.5014/ajot.2014.683001>
- Robertson, S. C. (1997). Mood disorders: Occupational therapy intervention. *OT Practice*, *2*(9), 34.
- Rogers, A. T., Bai, G., Lavin, R. A., & Anderson, G. F. (2016). Higher Hospital Spending on Occupational Therapy Is Associated With Lower Readmission Rates. *Medical Care Research and Review*. <https://doi.org/10.1177/1077558716666981>
- Rønningen, A., & Kjekken, I. (2008). Effect of an intensive hand exercise programme in patients with rheumatoid arthritis. *Scandinavian Journal of Occupational Therapy*, *15*, 173–183. (Level II)
- Ryan, S. (1995). Rheumatology. sharing care in an outpatient clinic. *Nursing Standard*, *10*(6), 23–25.
- Sackley, C., Wade, D. T., Mant, D., Atkinson, J. C., Yudkin, P., Cardoso, K., . . . Reel, K. (2006). Cluster randomized pilot controlled trial of an occupational therapy intervention for residents with stroke in UK care homes. *Stroke*, *37*, 2336–2341. (Level I)
- Salkeld, G., Cumming, R., O'Neill, E., Thomas, M., Szonyi, G., & Westbury, C. (2000). The cost effectiveness of a home hazard reduction program to reduce falls among older persons. *Australian and New Zealand Journal of Public Health*, *24*(3), 265–271. (Level I)
- Samėnienė, J., Kriščiūnas, A., & Endzelytė, E. (2008). The evaluation of the rehabilitation effects on cognitive dysfunction and changes in psychomotor reactions in stroke patients. *Medicina (Kaunas)*, *44*, 860–870. (Level III)
- Samnani, M. (2004). Passive exercises coupled with therapeutic activities: A comparative study in the management of frozen shoulder. *Indian Journal of Occupational Therapy*, *36*(2), 37–40. (Level III)
- Sandell, C. (2008). A multidisciplinary assessment and intervention for patients awaiting total hip replacement to improve their quality of life. *Journal of Orthopaedic Nursing*, *12*(1), 26–34.
- Schene, A. H., Koeter, M. W. J., Kikkert, M. J., Swinkels, J. A., & McCrone, P. (2007). Adjuvant occupational therapy for work related major depression works: Randomized trial including economic evaluation. *Psychological Medicine*, *37*, 351–362. (Level I)
- Schneider, J., Duggan, S., Cordingley, L., Mozley, C. G., & Hart, C. (2007). Costs of occupational therapy in residential homes and its impact on service use. *Aging & Mental Health*, *11*(1), 108–114. (Level II)
- Sewell, L., Singh, S. J., Williams, J. E. A., Collier, R., & Morgan, M. D. L. (2005). Can individualized rehabilitation improve function independence in elderly patients with COPD? *Chest*, *128*, 1194–1200. (Level I)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Sewell, L., Singh, S. J., Williams, J. E. A., Collier, R., & Morgan, M. (2006). How long should outpatient pulmonary rehabilitation be? A randomized controlled trial of four weeks versus seven weeks. *Thorax*, *61*, 767–771. (Level I)
- Shepherd, J., & Wilding, C. (2006). Occupational therapy for people with ventricular assist devices. *Australian Occupational Therapy Journal*, *53*(1), 47-49.
- Shimada, Y., Chida, S., Matsunaga, T., Sato, M., Hatakeyama, K., & Itoi, E. (2007). Clinical results of rehabilitation for accessory nerve palsy after radical neck dissection. *Acta Oto-Laryngologica*, *127*(5), 491-497.
- Siggeirsdottir, K., Olafsson, O., Jonsson, H., Iwarsson, S., Gudnason, V., & Jonsson, B. Y. (2005). Short hospital stay augmented with education and home-based rehabilitation improves function and quality of life after hip replacement. *Acta Orthopaedica*, *76*, 555–562. (Level I)
- Silva, A. C., Jones, A., Silver, P. G., & Natour, J. (2008). Effectiveness of a night-time hand positioning splint in rheumatoid arthritis: A randomized controlled trial. *Journal of Rehabilitation Medicine*, *40*, 749–754. (Level I)
- Singh, N., & Desai, O. P. (2007–2008). Prevention and management of low backache in pregnant women through the use of exercise program and education booklet. *Indian Journal of Occupational Therapy*, *39*(3), 65–72. (Level III)
- Slade, A., Tennant, A., & Chamberlain, M. A. (2002). A randomized controlled trial to determine the effect of intensity of therapy upon length of stay in a neurological rehabilitation setting. *Journal of Rehabilitation Medicine*, *34*, 260–266. (Level I)
- Sloane, P. D., Williams, C. D., Mitchell, C. M., Preisser, J. S., Wood, W., Barrick, A. L., . . . Zimmerman, S. (2007). High intensity environmental light in dementia: Effect on sleep and activity. *Journal of the American Geriatrics Society*, *55*, 1524–1533. (Level III)
- Smith, C., & Hale, L. (2007). The effects of non-pharmacological interventions on fatigue in four chronic illness conditions: A critical review. *Physical Therapy Reviews*, *12*(4), 324-334.
- Smith, M. (2008). What occupational therapy interventions have an effect on hand function in clients with rheumatoid arthritis? A systematic review. *Irish Journal of Occupational Therapy*, *36*(1), 31–38. (Systematic review)
- Somasundaram, S., & Sood, G. (2009). The effectiveness of task oriented group in reducing symptoms of post traumatic stress among females diagnosed with cancer of cervix. *Indian Journal of Occupational Therapy*, *41*(1), 9–13. (Level II)
- Sood, J. R., Cisek, E., Zimmerman, J., Zaleski, E. H., & Fillmore, H. H. (2003). Treatment of depressive symptoms during short-term rehabilitation: An attempted replication of the DOUR project. *Rehabilitation Psychology*, *48*(1), 44-49.
- Spencer, J., Hersch, G., Eschenfelder, V., Fournet, J., & Murray-Gerzik, M. (1999). Outcomes of protocol-based and adaptation-based occupational therapy interventions for low-income elderly persons on a transitional unit. *American Journal of Occupational Therapy*, *53*(2), 159-170.
- Spicka, C., Macleod, C., Adams, J., & Metcalf, C. (2009). Effect of silver ring splints on hand dexterity and grip strength in patients with rheumatoid arthritis: An observational pilot study. *Hand Therapy*, *14*(2), 53–57. (Level III)
- Stein, F., & Smith, J. (1989). Short-term stress management programme with acutely depressed in-patients. *Canadian Journal of Occupational Therapy*, *56*(4), 185-192.
- Steinman, L. E., Frederick, J. T., Prohaska, T., Satariano, W. A., Dornberg-Lee, S., Fisher, R., et al. (2007). Recommendations for treating depression in community-based older adults. *American Journal of Preventive Medicine*, *33*(3), 175-181.
- Stultjens, E. M. J., Dekker, J., Bouter, L. M., Leamrijse, C. J., & Van den Ende, C. H. M. (2005). Evidence of the efficacy of occupational therapy in different conditions: An overview of systematic reviews. *Clinical Rehabilitation*, *19*, 247–254. (Systematic review)

- Steultjens, E. M. J., Dekker, J., Bouter, L. M., van de Nes, J. C. M., Cup, E. H. C., & Van den Ende, C. H. M. (2003). Occupational therapy for stroke patients: A systematic review. *Stroke*, *34*, 676–687. (Systematic review)
- Steultjens, E. M., Dekker, J., Bouter, L. M., Cardel, M., Van de Nes, J. C., & Van den Ende, C. H. (2003). Occupational therapy for multiple sclerosis. *Cochrane Database Systemic Reviews*, Issue 3, Article No. CD003608. Reprinted in 2009. (Systematic review)
- Steultjens, E. M., Dekker, J., Bouter, L. M., Jellema, S., Bakker, E. B., & Van den Ende, C. H. (2004). Occupational therapy for community dwelling elderly people: A systematic review. *Age and Ageing*, *33*, 453–460. (Systematic review)
- Steultjens, E. M., Dekker, J., Bouter, L. M., Schaardenberg, D., Kuyk, M. A. H., & Van den Ende, C. H. (2004). Occupational therapy for rheumatoid arthritis. *Cochrane Database of Systematic Reviews*, Issue 1, Article No. CD003114 (Systematic review)
- Stevens, C. (1996). Complementary therapies in cancer care: An NHS approach. *International Journal of Palliative Nursing*, *2*(1), 15-18.
- Stitik, T. P., Foye, P. M., & Nadler, S. F. (1999). Osteoarthritis of the knee and hip: Part 2 -- keys to successful nondrug therapy. *Consultant (00107069)*, *39*(7), 1975.
- Stitik, T. P., Yonclas, P., Foye, P. M., & Schoenherr, L. (2005)a. Non pharmacologic management of knee and hip osteoarthritis. *Journal of Musculoskeletal Medicine*, *22*(2), 61.
- Stitik, T. P., Yonclas, P., Foye, P. M., & Schoenherr, L. (2005)b. Osteoarthritis of the knee and hip: Practical nondrug steps to successful therapy. *Consultant*, *45*(12), 1248-1258.
- Studenski, S., Duncan, P. W., Perera, S., Reker, D., Lai, S. M., & Richards, L. (2005). Daily functioning and quality of life in a randomized controlled trial of therapeutic exercise for subacute stroke survivors. *Stroke*, *36*, 1764–1770. (Level I)
- Taylor, K., & Currow, D. (2003). A prospective study of patient identified unmet activity of daily living needs among cancer patients at a comprehensive cancer care centre. *Australian Occupational Therapy Journal*, *50*(2), 79-85.
- Taylor, R. R. (2004). Quality of life and symptom severity for individuals with chronic fatigue syndrome: Findings from a randomized clinical trial. *American Journal of Occupational Therapy*, *58*, 35–43. (Level I)
- Taylor, R. R., Jason, L. A., Shiraishi, Y., Schoeny, M. E., & Keller, J. (2006). Conservation of resources theory, perceived stress, and chronic fatigue syndrome: Outcomes of a consumer-driven rehabilitation program. *Rehabilitation Psychology*, *51*, 157–165. (Level I)
- Taylor, R., & Marley, A. (2006). An integrated model for community COPD care. *Australian Journal of Primary Health*, *12*(2), 45-50.
- Thomas, J. J. (1996). Comparison of patient education methods: Effects on knowledge of cardiac rehabilitation principles. *Occupational Therapy Journal of Research*, *16*(3), 166-178.
- Tijhuis, G. J., VlietVlieland, T. P., Zwinderman, A. H., & Hazes, J. M. (1998). A comparison of the futuro wrist orthosis with a synthetic thermoLyn orthosis: Utility and clinical effectiveness. *Arthritis Care and Research*, *11*(3), 217-222.
- Timmons, A., & MacDonald, E. (2008). 'Alchemy and magic': The experience of using clay for people with chronic illness and disability. *British Journal of Occupational Therapy*, *71*(3), 86-94.
- Towheed, T. E. (2005). Systematic review of therapies for osteoarthritis of the hand. *Osteoarthritis & Cartilage*, *13*(6), 455-462.
- Trombly, C. A., Radomski, M. V., Trexel, C., & Burnet-Smith, S. E. (2002). Occupational therapy and achievement of selfidentified goals by adults with acquired brain injury: Phase II. *American Journal of Occupational Therapy*, *56*, 489–498. (Level III, comparative study)
- Tsai, A. Y., Yang, M., Lan, C., & Chen, C. (2008). Evaluation of effect of cognitive intervention programs for the community dwelling elderly with subjective memory complaints. *International Journal of Geriatric Psychiatry*, *23*, 1172–1174. (Level II)

AITO Associazione Italiana dei Terapisti Occupazionali

Sede legale e recapito postale: Via Angelo Emo 144, 00136 Roma
Email / sito web: info@aito.it / www.aito.it

- Tsauo, J. Y., Chen, W. H., Liang, H. W., & Jang, Y. (2009). The effectiveness of a functional training programme for patients with chronic low back pain: A pilot study. *Disability and Rehabilitation, 31*, 1100–1106. (Level I)
- Van Heugten, C. M., Dekker, J., Deelman, B. G., Stehmann-Saris, J. C., & Kinebanian, A. (2000). Rehabilitation of stroke patients with apraxia: The role of additional cognitive and motor impairments. *Disability and Rehabilitation, 22*, 547–554. (Level II)
- Van Heugten, C. M., Dekker, J., Deelman, B. G., van Dijk, A. J., Stehmann-Saris, F. C., & Kinebanian, A. (2000). Measuring disabilities in stroke patients with apraxia: A validation study of an observational method. *Neuropsychological Rehabilitation, 10*, 401–414. (Level II)
- Vockins, H. (2004). Occupational therapy intervention with patients with breast cancer: A survey. *European Journal of Cancer Care, 13*(1), 45–52.
- Von Koch, L., de Pedro-Cuesta, J., Kostulas, V., Almazán, J., & Holmqvist, L. W. (2001). Randomized controlled trial of rehabilitation at home after stroke: One-year follow-up of patient outcome, resource use and cost. *Cerebrovascular Diseases, 12*, 131–138. (Level I)
- Von Koch, L., Holmqvist, L. H., Hottrich, A. W., Tham, K., & De Pedro-Cuesta, J. (2000). Rehabilitation at home after stroke: A descriptive study of an individualized intervention. *Clinical Rehabilitation, 14*, 574–583. (Level I)
- WaiShanLouie, S. (2004). The effects of guided imagery relaxation in people with COPD. *Occupational Therapy International, 11*(3), 145–159.
- Warsi, A., LaValley, M. P., Wang, P. S., Avorn, J., & Solomon, D. H. (2003). Arthritis self-management education programs: A meta-analysis of the effect on pain and disability. *Arthritis & Rheumatism, 48*(8), 2207–2213.
- Waters, D. (1995). Recovering from a depressive episode using the canadian occupational performance measure. *Canadian Journal of Occupational Therapy, 62*(5), 278–282.
- Werngren-Elgström, M., Iwarsson, S., Elmståhl, S., & Dehlin, O. (2005). ADL dependence and perceived health among elderly deaf sign-language users: A comparison with matched groups of elderly hearing people. *Scandinavian Journal of Occupational Therapy, 12*(2), 81–88. (Level II)
- Wilde, C. K., & Hall, J. A. (1995). Occupational therapy in cardiac rehabilitation: Resumption of daily life activities. *Physical Medicine and Rehabilitation Clinics of North America, 6*(2), 349–372.
- Wilkins, S., Jung, B., Wishart, L., Edwards, M., & Norton, S. G. (2003). The effectiveness of community-based occupational therapy education and functional training programs for older adults: A critical literature review. *Canadian Journal of Occupational Therapy - Revue Canadienne d'Ergotherapie, 70*(4), 214–225.
- Winkel, A., Langberg, H., & Wæhrens, E. E. (2015). Reablement in a community setting. *Disability and Rehabilitation, 37*(15), 1347–1352. <https://doi.org/10.3109/09638288.2014.963707>
- Wiskin, L. F. (1998). Cognitive-behavioral therapy: A psychoeducational treatment approach for the american worker with rheumatoid arthritis. *Work, 10*(1), 41–48.
- Wynn, N., & Wickham, J. (2009). Night-time positioning for children with postural needs: What is the evidence to inform best practice? *British Journal of Occupational Therapy, 72*, 543–550. (Systematic review)
- Yakobina, S., & Tallant, B. K. (1997). I came, I thought, I conquered: Cognitivebehaviour approach applied in occupational therapy for the treatment of depressed (dysthymic) females. *Occupational Therapy in Mental Health, 13*(4), 59–73.
- Yuen, H. K., Mitcham, M., & Morgan, L. (2006). Managing post-therapy fatigue for cancer survivors using energy conservation training. *Journal of Allied Health, 35*(2), 121E–139E. (Level I)

- Zastrow, R., Van Gilder, T., & Quadracci, L. J. (2010). An Employer-Directed Health Plan That Seeks To Reenergize Primary Care. *Health Affairs*, *29*(5), 976–978. <https://doi.org/10.1377/hlthaff.2010.0031>
- Zhang, L., Abreu, B. C., Gonzales, V., Huddleston, N., & Ottenbacher, K. (2002). The effect of predictable and unpredictable motor tasks on postural control after traumatic brain injury. *NeuroRehabilitation*, *17*, 225–230. (Level II)
- Zijlstra, T. R., HeijnsdijkRouwenhorst, L., & Rasker, J. J. (2004). Silver ring splints improve dexterity in patients with rheumatoid arthritis. *Arthritis & Rheumatism (Arthritis Care & Research)*, *51*(6), 947-951.